



TITLE:

On annulus twists (Intelligence of Low-dimensional Topology)

AUTHOR(S):

Abe, Tetsuya

CITATION:

Abe, Tetsuya. On annulus twists (Intelligence of Low-dimensional Topology). 数理解析研究所講究録 2016, 2004: 108-114: KJ00010275649.

ISSUE DATE:

2016-07

URL:

<http://hdl.handle.net/2433/231504>

RIGHT:

On annulus twists

Tetsuya Abe

Osaka City University Advanced Mathematical Institute

ABSTRACT. We survey some results about annulus twists which are related to Dehn surgery on knots, knot concordance, and 4-manifold theory.

1. INTRODUCTION

Lickorish [15] and Wallace [22] proved that every closed connected orientable 3-manifold can be obtained by Dehn surgery on some link in S^3 . In other words, every closed connected orientable 3-manifold is described by a framed link in S^3 . Our interest is in uniqueness of framed link descriptions of a given 3-manifold. A natural question is the following.

Question 1. If two framed links \mathcal{L} and \mathcal{L}' give the same 3-manifold, then are \mathcal{L} and \mathcal{L}' isotopic as framed links ?

It is well-known that the answer of Question 1 is NO. Indeed, for a given framed link \mathcal{L} and a $1/n$ -framed unknot \mathcal{O} , two framed links \mathcal{L} and $\mathcal{L} \sqcup \mathcal{O}$ give the same 3-manifold. A modified question is the following.

Question 2. If two framed knots \mathcal{K} and \mathcal{K}' give the same 3-manifold, then are \mathcal{K} and \mathcal{K}' isotopic as framed knots ?

The answer of Question 2 is again NO [16] (see also [8, 9, 17, 20]). The remaining questions are the following.

- (1) Under what conditions, are framed knot descriptions of a 3-manifold unique?
- (2) To what extent, are framed knot descriptions of a 3-manifold far from unique ?

For the question (1), for example, see [12, 13, 14, 18]. We concentrate on the question (2). More precisely, we consider Clark's problem in Kirby problem list [10]:

Problem 3.6(D). Fix an integer n . Is there a homology 3-sphere (or any 3-manifold) which can be obtained by n -surgery on an infinite number of distinct knots?

In [19], Osoinach solved Problem 3.6(D) for the case $n = 0$ by constructing knots using the method of twisting along an annulus, which we call an annulus twist. After Teragaito's work [21] (see also [7, 11]), Jong, Luecke, Osoinach, and the author [3] solved Problem 3.6(D) affirmatively, where they generalized annulus twists.

In [4], a 4-dimensional extension of Problem 3.6(D) was proposed as follows:

Problem 1. Let n be an integer. Find infinitely many mutually distinct knots K_1, K_2, \dots such that $X_{K_i}(n) \approx X_{K_j}(n)$ for each $i, j \in \mathbb{N}$.

Here $X_K(n)$ denotes the smooth 4-manifold obtained from the 4-ball B^4 by attaching a 2-handle along K with framing n , and the symbol \approx stands for a diffeomorphism. Due to Akbulut [5, 6], there exists a pair of distinct knots K_n and K'_n such that $X_{K_n}(n) \approx X_{K'_n}(n)$ for each $n \in \mathbb{Z}$, which is a partial answer to Problem 1. In [4], Jong, Omae, Takeuch, and the author solved Problem 1 for the case $n = 0, \pm 4$. In [3], Jong, Luecke, Osoinach, and the author also solved Problem 1 affirmatively.

2. OSOINACH'S RESULT

In this section, we recall Osoinach's result in [19]. Let K_n be the knots in Figure 1, which is isotopic to the knots in the page 731 in [19]. One of the main results in [19] is the following.

Theorem 2.1 (Osoinach [19]). *We have the following.*

- (1) *The 3-manifold obtained by 0-surgery of K_0 is toroidal.*
- (2) *The sequence $\{K_n\}$ contains infinitely many distinct hyperbolic knots.*
- (3) *$S_0^3(K_0) \approx S_0^3(K_1) \approx S_0^3(K_2) \approx S_0^3(K_3) \approx \dots$, where $S_n^3(K)$ denotes the 3-manifold obtained by n -surgery of a knot K in S^3 .*




FIGURE 1. The definition of the knots K_n .

Note that we can check that K_n and K_{-n} are isotopic, and Takioka proved that the knots K_n ($n \geq 0$) are mutually distinct by calculating the Gamma polynomial which is a specialization of the HOMFLYPT polynomial.

Let V be the solid torus standardly embedded in S^3 and V' the 3-manifold as in Figure 2. The main observation in [19] is the following.




FIGURE 2. The definitions of V and V' .

Lemma 2.2 (cf. Theorem 2.1 in [19]). *There exists a (natural) diffeomorphism*

$$\varphi_n : V' \longrightarrow V$$

such that $\varphi_n|_{\partial V'} = id$.

Remark 2.3. *Osoinach [19] considered the diffeomorphism φ_n^{-1} .*

Figure 2 explains a proof of (3) in Theorem 2.1. Note that, by Lemma 2.2, the picture on the bottom-left is diffeomorphic to $S^3_0(K_0)$.




FIGURE 3. A proof of (3) in Theorem 2.1.

3. DEHN SURGERY AND KNOT CONCORDANCE

We recall a terminology in knot concordance. Two knots K and K' are *concordant* if they cobound a properly embedded annulus in $S^3 \times I$. In this paper, we do NOT consider orientations of a given knot.

Dehn surgery on knots and knot concordance are closely related. A motivating question is the following.

Question 3.1 (A. Levine [24]). *If K is concordant to K' , then for all n , $S_n^3(K)$ is homology cobordant to $S_n^3(K')$. Is the converse true?*

The following conjecture is due to Akbulut and Kirby (see Problem 1.19 in the Kirby's problem list [10]).

Conjecture. If 0-framed surgeries on two knots give the same 3-manifold, then the knots are concordant.

Tagami and the author [2] proved that Akbulut-Kirby's conjecture is false if the slice-ribbon conjecture is true. Subsequently, Yasui [23] proved that Akbulut-Kirby's conjecture is false by constructing knots K and K' satisfying

- (1) $X_K(0)$ and $X_{K'}(0)$ are exotic (i.e. homeomorphic but non-diffeomorphic).
- (2) K and K' are not concordant.

Note that $X_K(0)$ and $X_{K'}(0)$ are related by a cork twist. For the details, see [23]. The remaining conjecture is the following.

Conjecture. Let K and K' be knots. If $X_K(0)$ and $X_{K'}(0)$ are diffeomorphic, then K and K' are concordant.




FIGURE 4. The definition of K_0 and K_1 .

Remark: Let K_0 and K_1 be the knots in Figure 4. By the result in [4], the 4-manifolds $X_{K_0}(0)$ and $X_{K_1}(0)$ are diffeomorphic. Furthermore, if the slice-ribbon conjecture is true, K_0 and K_1 are not concordant (see [2]).

Acknowledgments. The author was supported by JSPS KAK-ENHI Grant Number 16K17597.

REFERENCES

- [1] T. Abe and M. Tange, *A construction of slice knots via annulus twists*, (2016), accepted by The Michigan Mathematical Journal.
- [2] T. Abe and K. Tagami, *Fibered knots with the same 0-surgery and the slice-ribbon conjecture*, (2016), to appear in Math. Research Letters.
- [3] T. Abe, I. Jong, J. Luecke and J. Osoinach, *Infinitely many knots admitting the same integer surgery and a 4-dimensional extension*, Int. Math. Res. Not. IMRN. (2015), doi: 10.1093/imrn/rnv008.
- [4] T. Abe, I. Jong, Y. Omae and M. Takeuchi, *Annulus twist and diffeomorphic 4-manifolds*, Math. Proc. Cambridge Philos. Soc. **155** (2013), 219–235.
- [5] S. Akbulut, *Knots and exotic smooth structures on 4-manifolds*, J. Knot Theory Ramifications **2** (1993), no. 1, 1–10.
- [6] S. Akbulut, *On 2-dimensional homology classes of 4-manifolds*. Math. Proc. Camb. Phil. Soc. **82** (1977), no. 1, 99–106.
- [7] K. L. Baker, C. McA. Gordon, and J. Luecke, *Bridge number and integral Dehn surgery*, Algebraic & Geometric Topology **16** (2016) 1–40.
- [8] R. Brakes, *Manifolds with multiple knot-surgery descriptions*, Math. Proc. Camb. Phil. Soc. **87** (1980), no. 3, 443–448.
- [9] A. Kawachi, *Mutative hyperbolic homology 3-spheres with the same Floer homology*, Geom.Dedicata **61** (1996), no. 2, 205–217.
- [10] R. Kirby, *Problems in low-dimensional topology*. AMS/IP Stud. Adv. Math. 2(2), Geometric topology (Athens, GA, 1993), (Amer. Math. Soc. 1997).
- [11] R. Kouno, *3-manifolds with infinitely many knot surgery descriptions (in Japanese)*, Masters thesis, Nihon University (2002).
- [12] D. Gabai, *Foliations and the topology of 3-manifolds. III*, J. Differential Geom. **26**, No. 3 (1987), 479–536.
- [13] P. Ghiggini *Knot Floer Homology Detects Genus-One Fibred Knots*, American Journal of Mathematics **130**, No. 5 (2008), 1151–1169
- [14] M. Lackenby, *Dehn Surgery on Knots in 3-Manifolds*, Journal of the American Mathematical Society, No. 4 (1997), 835–864.
- [15] W. B. R. Lickorish, *A representation of orientable combinatorial 3-manifolds*, Ann. of Math **76** (1962), 531–538.
- [16] W. B. R. Lickorish, *Surgery on knots*, Proc. Amer. Math. Soc. **60** (1976), 296–298.
- [17] C. Livingston, *More 3-manifolds with multiple knot-surgery and branched-cover descriptions*, Math. Proc. Camb. Phil. Soc. **91** (1982), no. 3, 473–475.
- [18] Yi Ni and Zhongtao Wu, *Cosmetic surgeries on knots in S^3* , J. Reine Angew. Math., **706**, (2015), 1–17.
- [19] J. Osoinach, *Manifolds obtained by surgery on an infinite number of knots in S^3* , Topology **45** (2006), 725–733.
- [20] M. Teragaito, *Toroidal Dehn surgery on hyperbolic knots and hitting number*, Topology Appl. **157** (2010), no. 1, 269–273.
- [21] M. Teragaito, *A Seifert fibered manifold with infinitely many knot-surgery descriptions*, Int. Math. Res. Not. **2007**, no. 9, Art. ID rnm028, 16 pp.
- [22] A. Wallace, *Modifications and cobounding manifolds*, Can. J. Math. **12** (1960), 503–528.
- [23] K. Yasui *Corks, exotic 4-manifolds and knot concordance*, arXiv:1505.02551v3 [math.GT].

- [24] Problem list Banff 2016, “Synchronisation of smooth and topological 4-manifolds”, Banff International Research Station, February 2016.

Osaka City University Advanced Mathematical Institute
3-3-138 Sugimoto, Sumiyoshi-ku Osaka 558-8585 JAPAN

E-mail address: tabe@sci.osaka-cu.ac.jp

大阪市立大学・数学研究所 安部 哲哉